



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

- Callitriche heterophylla*, Ph., is common on Long Island.
Quercus palustris, Du Roi., is found at Manor, Long Island.
Cupressus thyoides, L., is common around River Head, where it takes the place of the red cedar.
Potamogeton hybridus, Mx. Wading River.
Potamogeton pusillus, L., var. *tenuissimus*. River Head.
Potamogeton perfoliatus, L. Northville and River Head.
Potamogeton gramineus, var. *graminaefolius*. River Head.
Spiranthes graminea, Lindl., var. *Walteri*. Wading River.
Liparis Læselii, Richard. Wading River.
Lachnanthes tinctoria, Ell., was found at Manorville this year by Hon. Isaac Coles, of Glen Cove.
Sporobolus serotinus, Gray. River Head.
Muhlenbergia sylvatica, T. & G. River Head.
Elymus Canadensis, L., is common on Long Island.
Panicum amarum, Ell., is common at Wading River, on the sound, and at River Head, on the bay.
Cystopteris fragilis, Bernh. In an old well at Rocky Point, and in a deep ravine at Wading River; not a dozen plants in both localities.

E. S. MILLER.

Additions to the Flora of Onondaga County, N. Y.—In spite of very adverse weather the Syracuse Botanical Club has made some botanical excursions during the past season, and very satisfactory ones too. We have found the following plants, which were entirely new to us:

Viola striata, Ait. (with cream-colored flowers); *Corydalis flavula*, Raf; *Atriplex hortensis*, L., found by Mrs. Charles Barnes by the road near High Bridge; *Aplectrum hyemale*, Nutt. (with bright lemon-colored flowers); sterile plants of *Humulus Lupulus*, L.; *Polygonum Virginianum*, L.; *Quercus macrocarpa*, Mx.; *Lappa officinalis*, Allioni, (with white flowers); *Erythræa Centaurium*, Pers.; a sedge not identified; *Aster linifolius*, L.; *Aster puniceus*, L. (with rose-colored flowers); *Aster simplex*, Willd.; *Aster puniceus*, L., var. *vimineus*, Gray; *Helianthus decapetalus*, L.; *Asclepias phytolaccoides*, Pursh; and *Cladium mariscoides*, Torr. Mr. Beauchamp sent to us from Baldwinsville, *Spergularia rubra*, Presl., var. *campestris*, Gray. Besides the above we have collected many species not represented in our county herbarium.

Syracuse, N. Y.

MARY OLIVIA RUST.

Flora of Sam's Point.—The note by Mr. Britton in the last number of this journal, describing the botanical characteristics of Sam's Point, omits mention of some peculiarities which I noticed during a visit to this lovely spot in a most beautiful country, about the middle of September of the present year. I there saw, for the first time, the American mountain-ash (*Pyrus arbutifolia*), and was impressed with its splendid appearance. The European species (*P. aucuparia*) is cultivated in Washington and vicinity, and I have often admired the beautiful orange-colored berries; but our native species far sur-